

PYRIDINIUM AND QUINOLINIUM DERIVATIVES**DESCRIPTION****Field of the Invention**

The invention generally relates to compounds blocking
5 phosphorylcholine biosynthesis by means of the selective
blocking of the choline kinase enzyme in tumor cells or in cells
affected by parasitic infection and therefore being applicable
in the treatment of tumors and parasitic diseases or diseases
produced by viruses, bacteria and fungi in animals including
10 human beings; as well as to a method for preparing the compounds
of the invention and certain intermediates of said method.

Background of the Invention

Choline kinase is the first enzyme in the Kennedy or
phosphatidylcholine (PC) synthesis pathway and phosphorylates
15 choline to phosphorylcholine (PCho) using adenosine 5'-
triphosphate (ATP) as a phosphate group donor [Kent, C. *Prog.*
Lipid Res., 29, 87-105 (1990); Kennedy, E. P. *Fed. Proc.*, 20,
934-940 (1961)]. Ras genes form a family called oncogenes, which
have been widely studied because they are activated in 25-30% of
20 all human tumors and in some of them in 90% [Bos, J.L. *Cancer Res*
49, 4682-4689 (1989); Kiaris, H., Spandidos, D. A. *Int. J.*
Oncol., 413-421 (1995)]. Ras proteins play an important role in
intracellular signal transmission due to their involvement in
regulating cell proliferation, terminal differentiation and
25 senescence [Abdellatif, M., MacLellan, W. R.; Schneider, M. D.
J. Biol. Chem., 269, 15423-15426 (1994); Wiesmüller, L.,
Wittinghofer, F. *Cell Signal.*, 6, 247-267 (1994); Barbacid, M.
Eur. J. Clin. Invest., 20, 225-235 (1990); Hahn & Weinberg *Nat.*
Rev. Cancer, 2: 331 (2002); Wright & Shay *Nat. Biotech*, 20: 682
30 (2002); Drayton & Peters *Curr. Op. Gen. Dev*, 12:98 (2002)]. The
transformation mediated by various oncogenes, amongst which ras
oncogenes stand out, induces high levels of choline kinase
activity, resulting in an abnormal increase in the intracellular
levels of its product, PCho [Lacal et al., *Nature* 330, 269-272
35 (1987); Lacal J.C. *Mol. Cell. Biol.* 10, 333-340 (1990);
Teegarden, D., Taparowsky, E. J., Kent, C. *J. Biol. Chem.* 265,
6042-6047 (1990); Ratnam, S.; Kent, C. *Arch. Biochem. Biophys.*

323, 313-322 (1995); Ramírez de Molina, A., Rodríguez-González, A., Peñalva, V., Lucas, L., Lacal, J. C. *Biochem. Biophys. Res. Commun.* **285**, 873-879 (2001); Ramírez de Molina, A., Peñalva, V.; Lucas, L., Lacal, J. C. *Oncogene* **21**, 937-946 (2002)].

5 Supplementary facts support the role of ChoK in the generation of human tumors because studies using nuclear magnetic resonance (NMR) techniques have shown high levels of PCho in human tumor tissues with respect to normal tissues, including, among others, breast, colon, lung and prostate tumors [Ruiz-Cabello, J.,

10 Cohen, J. S. *NMR Biomed.* **5**, 226-233 (1992); de Certaines, J. D., Larsen, V. A., Podo, F., Carpinelli, G., Briot, O., Henriksen, O. *NMR Biomed.* **6**, 345-365 (1993); Smith, T. A. D., Bush, C., Jameson, C., Titley, J. C., Leach, M. O., Wilman, D. E. V., McCready, V. R. *NMR Biomed.* **6**, 318-323 (1993)]. It is common

15 knowledge that ras is one of the most profoundly studied oncogenes in human carcinogenesis and that ChoK inhibition has shown to be a new and efficient antitumor strategy in cells transformed by oncogenes [Cuadrado, A., Carnero, A., Dolfi, F., Jiménez, B., Lacal, J. C. *Oncogene*, **8**, 2959-2968 (1993);

20 Jiménez, B., del Peso, L., Montaner, S., Esteve, P. Lacal, J. C. *J. Cell Biochem.*, **57**, 141-149 (1995); Hernández-Alcoceba, R., Saniger, L., Campos, J., Núñez, M. C., Khaless, F., Gallo, M. Á., Espinosa, A., Lacal, J. C. *Oncogene*, **15**, 2289-2301 (1997)]. These first observations were later extrapolated *in vivo* in nude

25 mice [Hernández-Alcoceba, R., Fernández, F., Lacal, J. C. *Cancer Res.* **59**, 3112-3118 (1999)]. The research on ChoK inhibitors has identified Hemicholinium-3 (HC-3) as a relatively powerful and selective blocking agent [Cuadrado A., Carnero A., Dolfi F., Jiménez B. and Lacal J.C. *Oncogene* **8**, 2959-2968 (1993); Jiménez

30 B., del Peso L., Montaner S., Esteve P. and Lacal J.C. *J. Cell Biochem.* **57**, 141-149 (1995); Hernández-Alcoceba, R., Saniger, L., Campos, J., Núñez, M. C., Khaless, F., Gallo, M. Á., Espinosa, A., Lacal, J. C. *Oncogene*, **15**, 2289-2301 (1997)]. This choline homologue with a biphenyl structure has been used to

35 design new antitumor drugs. Since HC-3 is a powerful respiratory paralyzing agent, it is not a good candidate for its clinical use. The synthesis of several derivatives has been based on

structural modifications of HC-3 improving the ChoK inhibitory activity and suppressing its toxic effects. The inhibitory effect produced by bisquaternized symmetrical compounds on proliferation has been correlated with the ability to induce PCho production in whole cells [Hernández-Alcoceba, R., Saniger, L., Campos, J., Núñez, M. C., Khaless, F., Gallo, M. Á., Espinosa, A., Lacal, J. C. *Oncogene*, **15**, 2289-2301 (1997) and ES 2 117 950]. When the 1,2-ethylene-p-(bibenzyl dimethyl-diyl) residue was used as spacer between the two cationic pyridinium heads substituted in position 4 [Campos, J., Núñez, M. C., Rodríguez, V., Gallo, M. Á., Espinosa, A. *Bioorg. & Med. Chem. Lett.* **10**, 767-770 (2000)], the structures were evaluated by their ability to inhibit the isolated ChoK (in ex vivo conditions) [Lacal J.C. *IDrugs* **4**: 419-426 (2001)]. The 4-NR₂ group provided a considerable contribution and it was proposed [Campos, J., Núñez, M. C., Rodríguez, V., Gallo, M. Á., Espinosa, A. *Bioorg. & Med. Chem. Lett.* **10**, 767-770 (2000)] that the role of this group is electronic, by delocalization of the positive charge. The increase in ChoK activity in various human breast carcinomas has been published [Ramírez de Molina, A., Gutiérrez, R., Ramos, M. A., Silva, J. M., Silva, J., Sánchez, J. J., Bonilla, F., Lacal, J. C. *Oncogene* **21**, 4317-4322 (2002)]. It has recently been reported that ChoK alteration is a frequent event in some human tumors such as lung, colorectal and prostate tumors [Ramírez de Molina, A., Rodríguez-González, A., Gutiérrez, R., Martínez-Piñero, L., Sánchez, J. J., Bonilla, F., Rosell, R., Lacal, J. C. *Biochem. Biophys. Res. Commun.* **296**, 580-583 (2002)].


The bisquaternized pyridinium derivatives described in the state of the art and particularly in patent ES 2 117 950, show, however, high levels of toxicity, limiting their extended therapeutic application.

Therefore, in the state of the art there is a need to develop compounds having an activity blocking phosphorylcholine biosynthesis in tumor cells or in processes produced by parasitic, viral, bacterial or fungal infection, and, at the same time, having low levels of toxicity.

The authors of the present invention have discovered, after diligent research, that certain modifications in the structure of the compounds described in the state of the art and particularly in patent ES 2 117 950, have an unexpectedly and therefore surprisingly significant decrease in the levels of toxicity of said compounds of the state of the art.

Brief Description of the Invention

Therefore, the invención provides as its first object a family of compounds having formula I,



I

10


the structure of which is characterized by having two N-aryl-aminopyridinium groups joined by a spacer. In addition to acting as blocking agents of phosphorylcholine biosynthesis, by means of selectively blocking the choline kinase enzyme in tumor cells or potentially in processes produced by parasitic, viral, bacterial or fungal infections, the compounds of this family have low levels of toxicity.

In a second object, the invention provides the use of the compounds of formula I in medicine.

20 An additional object of the present invention consists of providing pharmaceutical formulations comprising at least one compound of formula I.


The invention provides, in another object, a method for preparing the compounds of formula I.

25 Finally, the invention provides the compounds of formula VII participating as starting compounds in the method for preparing compounds of formula I.



Detailed Description of the Invention

In its first object, the invention provides a family of
5 compounds corresponding to general formula I:




where

- Q^- represents the conjugate base of a pharmaceutically suitable organic or inorganic acid;
- 10 R_1 and R'_1 represent, independently of each other, a radical selected from the group formed by H and C_{1-6} alkyl optionally substituted by trifluoromethyl, hydroxyl or alkoxy;
- 15 R_2 and R'_2 represent, independently of each other, an aryl radical optionally substituted by halogen, trifluoromethyl, hydroxyl, C_{1-6} alkyl, amino or alkoxy;
- 20 R_3 and R'_3 represent, independently of each other, either a radical selected from the group formed by H, halogen, trifluoromethyl, hydroxyl, amino, alkoxy and C_{1-6} alkyl optionally substituted by trifluoromethyl, hydroxyl, amino or alkoxy, or together with R_4 and R'_4 respectively, and independently of each other, a -CH=CH-CH=CH- radical optionally substituted by
- 25 halogen, trifluoromethyl, hydroxyl, C_{1-6} alkyl, amino or alkoxy;
- R_4 and R'_4 represent, independently of each other, either a

radical selected from the group formed by H and C₁₋₆ alkyl optionally substituted by halogen, trifluoromethyl, hydroxyl, amino or alkoxy, or together with R₃ and R'₃ respectively, and independently of each other, a -CH=CH-CH=CH- radical optionally substituted by halogen, trifluoromethyl, hydroxyl, C₁₋₆ alkyl, amino or alkoxy; and
 A represents a spacer group.


In addition to acting as blocking agents of phosphorylcholine biosynthesis by means of selectively blocking the choline kinase enzyme in tumor cells or in cells affected by parasitic infection, the compounds belonging to this family are characterized by having levels of toxicity lower than those of compounds of similar structure known in the state of the art. This feature of the compounds of the invention is shown in the examples given below.

In view of the present invention, spacer group "A" is understood as any divalent organic structure acting as a joining link between the two pyridinium groups present in the structure defined by formula I. In a particular embodiment of the invention, the spacer A has a structure according to one of the formulas II, III, IV, V and VI. These formulas represent radicals; in them, the line — at the ends represents a bond, and not a methyl group.




II


where m, n and p represent integers which can have the following values: m = 0, 1; n = 0, 1-10; p = 0, 1; with the condition that m, n and p do not take the value of zero at the same time.




III



IV



V



VI

10 According to the present invention, the radicals R_1 and R'_1 , R_2 and R'_2 , as well as R_3 and R_4 , R'_3 and R'_4 can represent different radicals or the same radicals, giving rise to asymmetric or symmetric compounds.

15 In a particular embodiment of the invention, the radicals R_2 and R'_2 represent, independently of each other, a phenyl radical optionally substituted by halogen, trifluoromethyl, hydroxyl, C_{1-6} alkyl, amino and alkoxy. In another particular embodiment of the invention, the radicals R_1 and R'_1 represent a methyl radical, whereas the radicals R_2 and R'_2 represent
20 independently of each other a phenyl radical optionally substituted by one or more halogen substituents. In a third particular embodiment, both the radicals R_3 and R_4 and the radicals R'_3 and R'_4 together represent, although independently of each other, a $-CH=CH-CH=CH-$ radical optionally substituted by
25 one or more halogen substituents.

 The preferred compounds of the invention are shown in the following table I:

8
























Table I

N°	R ₃ , R ₄ *	NR ₁ R ₂	A	Code
1	H, H			ACG560B
2	H, H			ACG416B
3	H, H			ACG548B
4	H, H			ACG604A
5	-(CH=CH) ₂ -			RSM964A
6	-C ⁵ H=C ⁶ H- C ⁷ Cl=C ⁸ H-			RSM820C
7	-(CH=CH) ₂ -			RSM932A
8	C ⁵ H=C ⁶ H- C ⁷ Cl=C ⁸ H-			RSM824B
9	-(CH=CH) ₂ -			RSM936A
10	-C ⁵ H=C ⁶ H- C ⁷ Cl=C ⁸ H-			RSM828B

*R₃ and R₄ can mean either each one is hydrogen or both form a single radical.

5 Finally, in a preferred embodiment of the invention, the conjugate base of pharmaceutically suitable organic or inorganic acid Q represents Br (bromide) or F₆P (hexafluorophosphate).

The compounds of the invention have a selective effect on signaling pathways necessary for transforming certain oncogenes
 10 which do not affect normal cells with the same intensity and therefore, leave a sufficient margin for greater efficacy in

antitumor treatment.

On the other hand, the biological assays carried out by the authors of the invention allow extending this type of activity to antiviral, antiparasitic or antifungal activity because it is known that several parasites like *Plasmodium falciparum* or *Trypanosoma cruzi*, several viruses such as adenovirus, bacteria such as *Streptococcus pneumoniae* and fungi like *Candida albicans* require the metabolic pathway of phosphatidylcholine synthesis through choline kinase in order to complete their infective cycles in humans and animals. In this sense, the background in literature supports the role of ChoK in the intracellular metabolism in certain nucleosides in Hep-G2 cells [Martin, L. T.; Faraj, A.; Schinazi, R. F.; Gosselin, G.; Mathe, C.; Imbach, J.-L.; Sommadossi, J.-P. *Biochemical Pharmacology*, **53**, 75-87 (1997)], the use of ChoK as an enzymatic marker in parasitic diseases [Wunderlich, F.; Helwig, M.; Schillinger, G.; Vial, H.; Philippot, J.; Speth, V. *Molecular and Biochemical Parasitology*, **23**, 103-115 (1987); Ancelin, M. L.; Vial, H. J. *Biochimica et Biophysica Acta (BBA)- Lipids and Lipid Metabolism*, **875**, 52-58 (1986)], and the participation of ChoK in the biosynthesis of important phospholipids in viruses [Balakivera L., Schoen G., Thouvenin E., Chroboczek J. *J. Virol.* **77**:4858-4866 (2003)], bacteria [Whiting GC, Gillespie SH. *FEMS Microbiol Lett.* **138**:141-145 (1996)] and fungi [Mago N, Khuller GK. *J Med Vet Mycol.* **28**:355-362 (1990)]; Mago N, Khuller GK. *J Med Vet Mycol.* **28**:355-362 (1990)]. All these studies support that ChoK inhibition could have important therapeutic consequences in curing the diseases mentioned hereinbefore.

Therefore, in a second object, the invention provides the use of the compounds of formula I in medicine. Specifically, the compounds of formula I are claimed for their use in medicine. In a particular embodiment, the invention provides the compounds of formula I for the treatment of cancer, preferably, breast, lung, colorectal and pancreatic cancer. In another particular embodiment, the invention provides the compounds of formula I for the treatment of viral disease, preferably those caused by Adenovirus; as well as for antiparasitic treatment, preferably

for those diseases caused by *Plasmodium* or *Trypanosoma*; antibacterial treatment, preferably for those diseases caused by *Streptococcus*; and antifungal treatment, preferably for those diseases caused by *Candida*.

5 On the other hand, the use of a compound of formula I is claimed in the manufacture of a medicament. In a particular embodiment, the compound of formula I is used in the manufacture of a medicament for cancer, preferably breast, lung, colorectal or pancreatic cancer. In another particular embodiment, the
10 compound of formula I is used in the manufacture of a medicament for the treatment of viral diseases, preferably those caused by Adenovirus; as well as in the manufacture of a medicament for antiparasitic treatment, preferably for those diseases caused by *Plasmodium* or *Trypanosoma*; in the manufacture of a medicament
15 for the treatment of bacterial disease, preferably those caused by *Streptococcus*, and in the manufacture of a drug for the treatment of fungal diseases, preferably those caused by *Candida*.

 In its third object, the invention provides pharmaceutical
20 formulations comprising at least one compound of formula I as an active ingredient. Said pharmaceutical formulations can contain one or more excipients and/or carrier substances. Furthermore, said formulations can contain any other active ingredient inhibiting the function of the choline kinase enzyme.

25 The excipients, carrier substances and auxiliary substances must be pharmaceutically and pharmacologically tolerable, such that they can be combined with other components of the formulation or preparation and do not have adverse effects in the organism treated. The pharmaceutical compositions or
30 formulations include those that are suitable for oral or parenteral administration (including subcutaneous, intradermal, intramuscular and intravenous administration), although the best administration route depends on the conditions of the patient. The formulations can be in the form of single doses. the
35 formulations can be prepared according to methods known in the field of pharmacology. The amounts of active substances to be administered can vary according to the particularities of the


therapy.

The invention also provides a method for preparing the compounds of formula I. This object of the invention has two different embodiments according to whether the compound of formula I has the same or different aminopyridinium groups.

a) Process for obtaining the compounds of formula I in which the aminopyridinium groups are the same: The process comprises reacting the corresponding heterocyclic derivative of formula VII and the dihalogenated derivative AX_2 (where X represents the halogen atom: Cl, Br or I) in 2:1 molar amounts in an organic solvent. The reaction preferably takes place in butanone, in a closed tube and at a temperature of 90 to 110°C.

b) Process for obtaining the compounds of formula I in which the aminopyridinium groups are different: The process comprises reacting the corresponding heterocyclic derivative of formula VII and the dihalogenated derivative AX_2 (where X represents the halogen atom: Cl, Br or I) in 1:1 molar amounts in an organic solvent, in order to give a monoquaternized product which is again reacted with another different heterocyclic derivative molecule, in a 1:1 molar ratio, using another organic solvent that is more polar than the first one so that the previously formed monoquaternized salt can be dissolved. The first step of the reaction preferably takes place in butanone, in a closed tube and at a temperature of 90 to 110°C; whereas the second step is preferably carried out in ethanol in a closed tube and at a temperature of 90 to 110°C.

Finally, in its last object, the invention provides the compounds of formula VII participating as starting compounds in the method for preparing the compounds of formula I.




VII

where

R_1 represents a radical selected from the group formed

- by H and C₁₋₆ alkyl optionally substituted by trifluoromethyl, hydroxyl or alkoxy;
- R₂ represents an aryl radical optionally substituted by halogen, trifluoromethyl, hydroxyl, C₁₋₆ alkyl, amino or alkoxy;
- R₃ represents either a radical selected from the group formed by H, halogen, trifluoromethyl, hydroxyl, amino, alkoxy and C₁₋₆ alkyl optionally substituted by trifluoromethyl, hydroxyl, amino or alkoxy, or together with R₄ a -CH=CH-CH=CH- radical optionally substituted by halogen, trifluoromethyl, hydroxyl, C₁₋₆ alkyl, amino or alkoxy;
- R₄ represents either a radical selected from the group formed by H, and C₁₋₆ alkyl optionally substituted by halogen, trifluoromethyl, hydroxyl, amino or alkoxy, or together with R₃ a -CH=CH-CH=CH- radical optionally substituted by halogen, trifluoromethyl, hydroxyl, C₁₋₆ alkyl, amino or alkoxy.

Among the compounds preferred among the compounds of formula VII are the compounds of formula VIII:



Compound	R ¹	R ²	R
A	Me		H
B	Me		Cl

The following examples are set forth as an illustration of the present invention:

EXAMPLES

PREPARATION EXAMPLES

Compound 1 (code ACG560B): 1,1'-(benzene-1,3-diylmethylen)bis[4-(4-chloro-*N*-methylanilino)pyridinium] dibromide.

The mixture of 4-(4-chloro-*N*-methylaniline)pyridine (125 mg, 0.57 mmol) and 1,3-bis(bromomethyl)benzene (75 mg, 0.28 mmol) in dry butanone (40 ml) was heated in a closed tube at 100°C for 144 hours. After filtration and rigorous washing with butanone, EtOAc and Et₂O, compound 1 was obtained pure as a white solid (125.2 mg, 62.7%); m.p.: 197-198°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.48 (d, 4H, H-2,6_{pyr}, *J* = 6.6); 7.64 (d, 4H, H-3,5_{anil}, *J* = 8.6); 7.57 (s, 1H, H-2_{Ph}); 7.45 (d, 5H, H-2,6_{anil} and H-5_{Ph}; *J* = 8.6); 7.37 (d, 2H, H-4,6_{Ph}, *J* = 7.7); 6.95 (bs, 4H, H-3,5_{pyr}); 5.49 (s, 4H, CH₂N⁺); 3.46 (s, 6H, Me). ¹³C-NMR (100 MHz, DMSO-*d*₆) δ 156.20 (C-4_{pyr}); 142.75 (C-2,6_{pyr}); 141.96 (C-1_{anil}); 136.18 (C-1.3_{Ph}); 132.78 (C-4_{anil}); 130.50 (C-3,5_{anil}); 129.73 (C-5_{Ph}); 128.37 (C-2,6_{anil}); 128.18 (C-4,6_{Ph}); 127.89 (C-2_{Ph}); 109.15 (C-3,5_{pyr}); 59.16 (CH₂N⁺); 41.42 (Me). HRMS (*m/z*): Calculated for C₃₂H₃₀N₄Cl₂Br (M - Br)⁺ 619.1031; found: 619.1031. Analysis for C₃₂H₃₀N₄Cl₂Br₂·1H₂O. Calculated: C 53.43; H 4.56; N 7.63%. Found: C 53.14; H 4.48; N 7.79%.

Compound 2 (code ACG416B): 1,1'-(biphenyl-3,3'-diylmethylen)bis[4-(*N*-methylanilino)pyridinium] dibromide.

The mixture of 4-(*N*-methylaniline)pyridine (216 mg, 1.17 mmol) and 3,3'-bis(bromomethyl)biphenyl (200 mg, 0.58 mmol) in dry butanone (40 ml) was heated in a closed tube at a 100°C for 24 hours. After filtration and thorough washing with butanone, the solid product was purified by recrystallization from MeOH and the residue was ground with Et₂O. Compound 2 was obtained as a white solid (294 mg, 71.5%); m.p.: 124-125°C. ¹H-NMR (300 MHz, CD₃OD) δ 8.35 (bs, 4H, H-2,6_{pyr}); 7.84 (s, 2H, H-2_{Ph}); 7.67 (d, 2H, H-6_{Ph}, *J* = 7.7); 7.56 (t, 4H, H-3,5_{anil}, *J* = 7.6); 7.50-7.44 (m, 4H, H-5_{Ph} and H-4_{anil}); 7.39 (d, 2H, H-4_{Ph}, *J* = 7.7); 7.33 (d, 4H, H-2,6_{anil}, *J* = 7.5); 6.95 (bs, 4H, H-3,5_{pyr}); 5.47 (s, 4H, CH₂N⁺); 3.51 (s, 6H, Me). ¹³C-NMR (75 MHz, CD₃OD) δ 158.48 (C-4_{pyr}); 144.82 (C-1_{anil}); 143.80 (C-2.6_{pyr}); 142.60 (C-1_{Ph}); 136.82 (C-3_{Ph}); 132.01 (C-3,5_{anil}); 131.14 (C-5_{Ph}); 130.12 (C-4_{anil});

128.99 (C-4_{Ph}); 128.82 (C-6_{Ph}); 128.58 (C-2_{Ph}); 127.52 (C-2,6_{anil});
 110.29 (C-3,5_{pyr}); 61.97 (CH₂N⁺); 41.42 (Me). HRMS (m/z):
 Calculated for C₃₈H₃₆N₄Br (M - Br)⁺ 627.2123; found: 627.2122.
 Analysis for C₃₈H₃₆N₄Br₂·2.5H₂O. Calculated: C 60.56; H 5.48; N
 5 7.43%. Found: C 60.70; H 5.83; N 7.20%.

Compound 3 (code ACG548B): 1,1'-(biphenyl-3,3'-
 diylmethylen)bis[4-(4-chloro-N-methylanilino)pyridinium]
 dibromide.

The mixture of 4-(4-chloro-N-methylaniline)pyridine (235
 10 mg, 1.07 mmol) and 3,3'-bis(bromomethyl)biphenyl (183 mg, 0.53
 mmol) in dry butanone (40 ml) was heated in a closed tube at
 100°C for 24 hours. After filtration and thorough washing with
 CHCl₃, the solid product was purified by recrystallization from
 MeOH, after adding Et₂O until turbidity. Compound 3 was obtained
 15 as a white solid (205 mg, 49.7%); m.p.: 279-280°C. ¹H-NMR (300
 MHz, DMSO-d₆) δ 8.57 (d, 4H, H-2,6_{pyr}, J = 6.5); 7.88 (s, 2H, H-
 2_{Ph}); 7.67 (d, 2H, H-6_{Ph}, J = 7.7); 7.61 (d, 4H, H-3,5_{anil}, J =
 8.6); 7.51 (t, 2H, H-5_{Ph}, J = 7.7); 7.42 (d, 6H, H-4_{Ph} and H-
 2,6_{anil}, J = 8.6); 6.99 (bs, 4H, H-3,5_{pyr}); 5.51 (s, 4H, CH₂N⁺);
 20 3.43 (s, 6H, Me). ¹³C-NMR (75 MHz, DMSO-d₆) δ 156.20 (C-4_{pyr});
 142.72 (C-2,6_{pyr}); 142.05 (C-1_{anil}); 140.01 (C-1_{Ph}); 136.20 (C-3_{Ph});
 132.79 (C-4_{anil}); 130.53 (C-3,5_{anil}); 129.73 (C-5_{Ph}); 128.47 (C-
 2,6_{anil}); 127.51 (C-4_{Ph}); 127.14 (C-6_{Ph}); 127.04 (C-2_{Ph}); 109.20 (C-
 3,5_{pyr}); 59.55 (CH₂N⁺); 40.73 (Me). HRMS (m/z): Calculated for
 25 C₃₈H₃₄N₄Cl₂Br (M - Br)⁺ 695.1344; found: 695.1344. Analysis for
 C₃₈H₃₄N₄Cl₂Br₂·1.2H₂O. Calculated: C 57.12; H 4.59; N 7.01%. Found:
 C 57.55; H 4.99; N 6.97%.

Compound 4 (code ACG604A): 1,1'-(biphenyl-3,3'-
 diylmethylen)bis[4-(3,5-dichloro-N-methylanilino)pyridinium]
 30 dibromide.

The mixture of 4-(3,5-dichloro-N-methylaniline)pyridine
 (200 mg, 0.80 mmol) and 3,3'-bis(bromomethyl)biphenyl (136 mg,
 0.40 mmol) in dry butanone (40 ml) was heated in a closed tube
 at 100°C for 72 hours. After filtration and thorough washing
 35 with butanone Et₂O, compound 4 was obtained as a pure white
 solid (270 mg, 79.7%); m.p.: 312-313°C. ¹H-NMR (300 MHz, DMSO-d₆)

δ 8.63 (d, 4H, H-2,6_{pyr}, J = 7.1); 7.92 (s, 2H, H-2_{ph}); 7.75 (s, 2H, H-4_{anil}); 7.70 (d, 2H, H-6_{ph}, J = 7.6); 7.62 (d, 4H, H-2,6_{anil}, J = 1.8); 7.53 (t, 2H, H-5_{ph}, J = 7.6); 7.45 (d, 2H, H-4_{ph}, J = 7.6); 7.04 (d, 4H, H-3,5_{pyr}, J = 7.1); 5.56 (s, 4H, CH₂N⁺); 3.44 (s, 6H, Me). ¹³C-NMR (75 MHz, DMSO-*d*₆) δ 156.20 (C-4_{pyr}); 145.27 (C-1_{anil}); 142.86 (C-2,6_{pyr}); 140.08 (C-1_{ph}); 136.11 (C-3_{ph}); 135.34 (C-3,5_{anil}); 129.70 (C-5_{ph}); 128.33 (C-4_{anil}); 127.55 (C-4_{ph}); 127.14 (C-6_{ph}); 127.07 (C-2_{ph}); 125.97 (C-2,6_{anil}); 109.53 (C-3,5_{pyr}); 59.65 (CH₂N⁺); 40.59 (Me). HRMS (*m/z*): Calculated for C₃₈H₃₂N₄Cl₄Br (M - Br)⁺ 763.0564; found: 763.0563. Analysis for C₃₈H₃₂N₄Cl₄Br₂·0.1H₂O. Calculated: C 53.81; H 3.81; N 6.60%. Found: C 53.41; H 4.19; N 6.25%.

Compound 5 (code RSM964A): 1,1'-(biphenyl-3,3'-diylmethylen)bis[4-(4-chloro-*N*-methylanilino)quinolinium] dibromide.

The mixture of 4-(4-chloro-*N*-methylanilino)quinoline (212 mg, 0.78 mmol) and 3,3'-bis(bromomethyl)benzene (134 mg, 0.39 mmol) in dry butanone (40 ml) was heated in a closed tube at 100°C for 72 hours. After filtration and thorough washing with butanone, EtOAc and Et₂O, compound 5 was obtained as a pure yellowish solid (134 mg, 40%); m.p.: 217-218°C. ¹H-NMR (300 MHz, DMSO-*d*₆): δ 9.24 (d, J = 7.4, 2H, H-2_{quin}); 8.18 (d, J = 8.9, 2H, H-8_{quin}); 7.84 (s, 2H, H-2_{ph}); 7.63 (d, J = 7.5, 2H, H-5_{quin}); 7.56-7.43 (m, 18H, H-5,6_{ph}, H-2,3,5,6_{anil}, H-3,6,7_{quin}); 7.23 (d, J = 7.4, 2H, H-4_{ph}); 6.08 (s, 4H, N⁺-CH₂); 3.74 (s, 6H, Me). ¹³C-NMR (100 MHz, DMSO-*d*₆): δ 157.87 (C-4_{quin}); 147.46 (C-2_{quin}); 146.42 (C-1_{anil}); 140.03 (C-1_{ph}); 138.83 (C-8_{aquin}); 135.61 (C-3_{ph}); 133.50 (C-7_{quin}); 131.69 (C-4_{anil}); 130.27 (C-3,5_{anil}); 129.62 (C-5_{ph}); 127.35 (C-6_{ph}); 127.18 (C-2,6_{anil}); 126.73 (C-6_{quin}); 126.09 (C-4_{ph}); 125.87 (C-5_{quin}); 125.67 (C-2_{ph}); 119.65 (C-4_{aquin}); 119.14 (C-8_{quin}); 107.10 (C-3_{quin}); 57.28 (N⁺-CH₂); 44.94 (Me). HRMS (*m/z*): Calculated for C₄₆H₃₈N₄Cl₂Br₂ [(M - Br)]⁺ 795.1657. Found: 795.1656. Analysis for C₄₆H₃₈N₄Cl₂Br₂·3H₂O. Calculated: C 59.31; H 4.76; N 6.01%. Found: C 59.24; H 4.70; N 5.65%.

Compound 6 (code RSM820C): 1,1'-(biphenyl-3,3'-diylmethylen)bis[4-(4-chloro-*N*-methylanilino)-7-

chloroquinolinium] dibromide.

The mixture of 7-chloro-4-(4-chloro-*N*-methylanilino)quinoline (300 mg, 0.98 mmol) and 3,3'-bis(bromomethyl)biphenyl (168 mg, 0.49 mmol) in dry butanone (40 ml) was heated in a closed tube at 100°C for 72 hours. After filtration and thorough washing with butanone and CHCl₃, the solid product was purified by recrystallization from EtOH or EtOH/MeOH, after adding Et₂O until turbidity. Compound 6 was obtained as a yellowish solid (154 mg, 45%); m.p.: 220-221°C.

¹H-NMR (300 MHz, DMSO-*d*₆): δ 9.19 (d, *J* = 7.5, 2H, H-2_{quin}); 8.29 (d, *J* = 1.7, 2H, H-8_{quin}); 7.85 (s, 2H, H-2_{Ph}); 7.64 (d, *J* = 7.2, 2H, H-5_{quin}); 7.57-7.45 (m, 16H, H-5,6_{Ph}, H-2,3,5,6_{anil}, H-3,6_{quin}); 7.25 (d, *J* = 7.7, 2H, H-4_{Ph}); 6.08 (s, 4H, N⁺-CH₂); 3.73 (s, 6H, Me). ¹³C-NMR (75 MHz, DMSO-*d*₆): δ 157.68 (C-4_{quin}); 148.01 (C-2_{quin}); 146.14 (C-1_{anil}); 140.14 (C-1_{Ph}); 139.85 (C-8a_{quin}); 138.48 (C-7_{quin}); 135.51 (C-3_{Ph}); 132.11 (C-4_{anil}); 130.50 (C-3,5_{anil}); 129.80 (C-5_{Ph}); 129.45 (C-6_{Ph}); 127.32 (C-2,6_{anil}); 126.89 (C-6_{quin}); 126.12 (C-4_{Ph}); 125.91 (C-5_{quin}); 125.82 (C-2_{Ph}); 118.48 (C-8_{quin}); 118.35 (C-4a_{quin}); 107.38 (C-3_{quin}); 57.14 (N⁺-CH₂); 45.18 (Me). HRMS (*m/z*): Calculated for C₄₆H₃₆N₄Cl₄Br₂ [(M - HBr - Br)]⁺ 783.1616. Found: 783.1616. Analysis for C₄₆H₃₆N₄Cl₄Br₂·1.5H₂O. Calculated: C 56.76; H 4.04; N 5.76%. Found: C 56.72; H 4.18; N 5.71%.

Compound 7 (code RSM932A): 1,1'-(biphenyl-4,4'-diylmethylen)bis[4-(4-chloro-*N*-methylanilino)quinolinium] dibromide.

The mixture of 4-(4-chloro-*N*-methylaniline)quinoline (240 mg, 0.89 mmol) and 4,4'-bis(bromomethyl)biphenyl (152 mg, 0.44 mmol) in dry butanone (40 ml) was heated in a closed tube at 100°C for 72 hours. After filtration and thorough washing with butanone, compound 7 was obtained as a pure yellowish solid (121 mg, 30%); m.p.: 255-257°C. ¹H-NMR (300 MHz, DMSO-*d*₆): δ 9.19 (d, *J* = 7.4, 2H, H-2_{quin}); 8.12 (d, *J* = 8.9, 2H, H-8_{quin}); 7.83 (pst, *J* = 7.5, 2H, H-7_{quin}); 7.66 (d, *J* = 8.2, 2H, H-5_{quin}); 7.55 (d, *J* = 8.8, 4H, H-3,5_{anil}); 7.44 (d, *J* = 8.9, 4H, H-2,6_{anil}); 7.56-7.39 (m, 12H, H-2,3,5,6_{Ph}, H-3_{quin}, H-6_{quin}); 6.05 (s, 4H, N⁺-CH₂); 3.73 (s, 6H, Me). ¹³C-NMR (75 MHz, DMSO-*d*₆): δ 157.86 (C-4_{quin}); 147.41

(C-2_{quin}); 146.40 (C-1_{anil}); 139.11 (C-1_{ph}); 138.78 (C-8a_{quin});
 134.30 (C-4_{ph}); 133.47 (C-7_{quin}); 131.69 (C-4_{anil}); 130.26 (C-
 3,5_{anil}); 127.34 (C-3,5_{ph}); 127.18 (C-2,6_{anil}), (C-2,6_{ph}); 127.08
 (C-6_{quin}); 126.08 (C-5_{quin}); 119.65 (C-4a_{quin}); 119.12 (C-8_{quin});
 5 107.06 (C-3_{quin}); 56.94 (N⁺-CH₂); 44.94 (Me). HRMS (m/z):
 Calculated for C₄₆H₃₈N₄Cl₂Br₂ [(M - Br)]⁺ 795.1657. Found:
 795.1658. Analysis for C₄₆H₃₈N₄Cl₂Br₂·2H₂O. Calculated: C 60.48; H
 4.63; N 6.13%. Found: C 60.06; H 4.48; N 5.87%.

**Compound 8 (code RSM824B): 1,1'-(biphenyl-4,4'-
 10 diylmethene)bis[4-(4-chloro-N-methylanilino)-7-
 chloroquinolinium] dibromide.**

The mixture of 7-chloro-4-(4-chloro-N-methylanilino)quinoline (300 mg, 0.98 mmol) and 4,4'-bis(bromomethyl)biphenyl (168 mg, 0.49 mmol) in dry butanone
 15 (100 ml) was heated in a closed tube at 100°C for 72 hours. After filtration and thorough washing with butanone, compound 8 was obtained as a pure yellowish solid (195 mg, 48%); m.p.: 276-277°C. ¹H-NMR (400 MHz, DMSO-d₆): δ 9.14 (d, J = 7.4, 2H, H-2_{quin}); 8.23 (d, J = 1.6, 2H, H-8_{quin}); 7.73 (d, J = 8.3, 2H, H-5_{quin});
 20 7.69 (d, J = 8.4, 4H, H-2,6_{ph}); 7.56 (d, J = 8.8, 4H, H-3,5_{anil}); 7.46 (d, J = 8.9, 4H, H-2,6_{anil}); 7.50-7.46 (m, 6H, H-6_{quin}, H-3_{quin}); 7.41 (d, J = 8.4, 4H, H-3,5_{ph}); 6.04 (s, 4H, N⁺-CH₂); 3.73 (s, 6H, Me). ¹³C-NMR (100 MHz, DMSO-d₆): δ 157.69 (C-4_{quin}); 147.98 (C-2_{quin}); 146.13 (C-1_{anil}); 139.82 (C-8a_{quin}); 139.21
 25 (C-1_{ph}); 138.51 (C-7_{quin}); 134.22 (C-4_{ph}); 132.14 (C-4_{anil}); 130.50 (C-3,5_{anil}); 129.45 (C-2,6_{anil}); 127.54 (C-3,5_{ph}); 127.33 (C-6_{quin}); 127.23 (C-2,6_{ph}); 126.52 (C-5_{quin}); 118.47 (C-8_{quin}); 118.35 (C-4a_{quin}); 107.33 (C-3_{quin}); 56.83 (N⁺-CH₂); 45.19 (Me). HRMS (m/e):
 Calculated for C₄₆H₃₆N₄Cl₄Br₂ [(M-HBr-Br)]⁺ 783.1616. Found:
 30 783.1614. Analysis for C₄₆H₃₆N₄Cl₄Br₂. Calculated: C 58.38; H 3.83; N 5.92%. Found: C 58.73; H 3.96; N 5.74%.

Compound 9 (code RSM936A): 1,1'-[ethylenebis(benzene-1,4-diylmethene)]bis[4-(4-chloro-N-methylanilino)quinolinium] dibromide.

35 The mixture of 4-(4-chloro-N-methylanilino)quinoline (204 mg, 0.76 mmol) and 4,4'-bis(bromomethyl)bibenzyl (140 mg, 0.37 mmol) in dry butanone (40 ml) was heated in a closed tube at

100°C for 72 hours. After filtration and thorough washing with butanone and CHCl_3 , compound 9 was obtained as a pure yellowish solid (70 mg, 20%); m.p.: 212-214°C. $^1\text{H-NMR}$ (300 MHz, $\text{DMSO}-d_6$): δ 9.19 (d, $J = 7.4$, 2H, H-2_{quin}); 8.10 (d, $J = 8.9$, 2H, H-8_{quin}); 7.82 (pst, $J = 7.5$, 2H, H-7_{quin}); 7.54 (d, $J = 8.8$, 4H, H-3,5_{anil}); 7.44 (d, $J = 8.9$, 4H, H-2,6_{anil}); 7.52-7.39 (m, 6H, H-3_{quin}, H-5_{quin}, H-6_{quin}); 7.24 (s, 8H, H-2,3,5,6_{ph}); 5.98 (s, 4H, N^+-CH_2); 3.73 (s, 6H, Me); 2.80 (s, 4H, CH_2-Ph). $^{13}\text{C-NMR}$ (100 MHz, $\text{DMSO}-d_6$): δ 157.80 (C-4_{quin}); 147.34 (C-2_{quin}); 146.44 (C-1_{anil}); 141.55 (C-1_{ph}); 138.74 (C-8_{aquin}); 133.36 (C-7_{quin}); 132.32 (C-4_{ph}); 131.63 (C-4_{anil}); 130.25 (C-3,5_{anil}); 128.79 (C-3,5_{ph}); 127.26 (C-6_{quin}); 127.17 (C-2,6_{anil}); 126.74 (C-2,6_{ph}); 126.04 (C-5_{quin}); 119.66 (C-4_{aquin}); 119.19 (C-8_{quin}); 107.06 (C-3_{quin}); 57.10 (N^+-CH_2); 44.93 (Me); 36.22 (CH_2-Ph). HRMS (m/z): Calculated for $\text{C}_{48}\text{H}_{42}\text{N}_4\text{Cl}_2\text{Br}_2$ [(M - Br)]⁺ 823.1970. Found: 823.1970. Analysis for $\text{C}_{48}\text{H}_{42}\text{N}_4\text{Cl}_2\text{Br}_2 \cdot 1\text{H}_2\text{O}$. Calculated: C 62.42; H 4.80; N 6.07%. Found: C 62.29; H 4.59; N 6.09%.


Compound 10 (code RSM828B): 1,1'-[ethylenebis(benzene-1,4-diylmethylen)]bis[4-(4-chloro-N-methylanilino)-7-chloroquinolinium] dibromide.

The mixture of 7-chloro-4-(4-chloro-N-methylanilino)quinoline (300 mg, 0.98 mmol) and 4,4'-bis(bromomethyl)bibenzyl (182 mg, 0.49 mmol) in dry butanone (40 ml) was heated in a closed tube at 100°C for 72 hours. After filtration and thorough washing with butanone, compound 10 was obtained as a pure yellowish solid (229 mg, 48%); m.p.: 256-257°C. $^1\text{H-NMR}$ (400 MHz, $\text{DMSO}-d_6$): δ 9.11 (d, $J = 7.4$, 2H, H-2_{quin}); 8.18 (d, $J = 1.5$, 2H, H-8_{quin}); 7.55 (d, $J = 8.8$, 4H, H-3,5_{anil}); 7.46 (d, $J = 8.8$, 4H, H-2,6_{anil}); 7.56-7.44 (m, 6H, H-3_{quin}, H-5_{quin}, H-6_{quin}); 7.24 (s, 8H, H-2,3,5,6_{ph}); 5.97 (s, 4H, N^+-CH_2); 3.72 (s, 6H, Me); 2.82 (s, 4H, CH_2-Ph). $^{13}\text{C-NMR}$ (100 MHz, $\text{DMSO}-d_6$): δ 157.63 (C-4_{quin}); 147.91 (C-2_{quin}); 146.16 (C-1_{anil}); 141.74, 139.75 and 138.88 (C-7_{quin}, C-8_{aquin} and C-4_{ph}); 132.20 (C-4_{anil}); 132.08 (C-1_{ph}); 130.50 (C-3,5_{anil}); 129.39 (C-6_{quin}); 128.99 (C-3,5_{ph}); 127.32 (C-2,6_{anil}); 126.90 (C-2,6_{ph}); 126.48 (C-5_{quin}); 118.55 (C-8_{quin}); 118.35 (C-4_{aquin}); 107.32 (C-3_{quin}); 57.02 (N^+-CH_2); 45.17 (Me); 36.33 (CH_2-Ph). HRMS (m/z): Calculated for

$C_{48}H_{40}N_4Cl_4Br_2$ [(M-HBr-Br)]⁺ 811.1927. Found: 811.1926. Analysis for $C_{48}H_{40}N_4Cl_4Br_2 \cdot 2H_2O$. Calculated: C 57.05; H 4.39; N 5.54%. Found: C 57.14; H 4.07; N 5.46%.

REAGENT PREPARATION


- 5 The compound α, α' -dibromo-*m*-xylene is a commercial product supplied by Sigma-Aldrich Química S. A. with address at Avenida Valdelaparra No. 51-53, 28100 Alcobendas (Madrid).



α, α' -dibromo-*m*-xylene

- 10 The following starting materials were prepared by means of the methods described in the respective references


1.- 3,3'-bis(bromomethyl)biphenyl



Werner, W. *J. Org. Chem.* **17**, 523-528 (1952)

15


2.- 4,4'-bis(bromomethyl)biphenyl



Szendey, G. L., Munnes, S. *Chem. Ber.* **94**, 38-42 (1961); Staab, H. A., Haenel, M. *Chem. Ber.* **106**, 2190-2202 (1973)

20


3.- Bis-*p*-(bromomethyl)bibenzyl



Cram, D. J., Steinberg, J. *J. Am. Chem. Soc.* **73**, 5691-5704 (1951)

25


4.- 4-(*N*-Methylanilino)pyridine



Campos, J., Núñez, M. C., Sánchez, R., Gómez-Vidal, J. A., Rodríguez-González, A., Báñez, M., Gallo, M. Á., Lacal, J. C.,


Espinosa, A. *Bioorg. & Med. Chem.* **10**, 2215-2231 (2002)

5.- 4-(4-Chloro-*N*-methylanilino)pyridine



- 5 Conejo-García, A., Campos, J., Sánchez, R., Rodríguez-González, A., Lacal, J. C., Gallo, M. Á., Espinosa, A. *Eur. J. Med. Chem.* **38**, 109-116 (2003).

6.- 4-(3,5-dichloro-*N*-methylanilino)pyridine




10

- This compound was prepared from 4-chloropyridine hydrochloride and 4-(3,5-dichloro-*N*-methylanilino)pyridine according to the process described previously in: Conejo-García, A., Campos, J., Sánchez, R., Rodríguez-González, A., Lacal, J. C., Gallo, M. Á., Espinosa, A. *Eur. J. Med. Chem.* **38**, 109-116 (2003). On the other hand, 3,5-dichloro-*N*-methylanilino was obtained by following the process described in the following work: Leeson, P. D., Baker, R., Carling, R. W., Curtis, N. R., Moore, K. W., Williams, B. J., Foster, A. C., Donald, A. E., Kemp, J. A., Marshall, G. R. *J. Med. Chem.* **34**, 1243-1252 (1991).


20

7.- 4,4'-Bis(chloromethyl)-[2,2']bithiazolyl.




- 25 Ref.: Chi, AND. F.; Chu, T. I. *Record (Peking)*, **1**, 45 (1957); *Chem. Abstract*, **52**, 6321 a,b (1957).

8.- Diethyl 4,4'-Bis(bromomethyl)-[2,2']bithiazolyl-5,5'-dicarboxylate




Ref.: Lehn, J.-M.; Regnouf de Vains, J.-B. *Tetrahedron Lett.*, **30**, 2209-2212 (1989).

5 9.- 6,6'-Bis(bromomethyl)-[2,2']bipyridine



Ref.: Rodríguez-Ubis, J.-C.; Alpha, B.; Plancherel, D.; Lehn, J.-M. *Helv. Chim. Acta*, **67**, 2264 (1984).


10 10.- 6,6'-Bis(bromomethyl)-4,4'-dimethyl-[2,2']bipyrimidinyl



Ref.: Lehn, J.-M.; Regnouf de Vains, J.-B. *Tetrahedron Lett.*, **30**, 2209-2212 (1989).

PREPARATION OF NEW STARTING MATERIALS


15 The compounds of formula VII:



VII

can be prepared by reacting the 4-aniline or quinoline derivative with the corresponding 4-chloro-aniline in glacial acetic acid with reflux. After cooling, the solution is basified with sodium hydroxide solution and the resulting suspension is subsequently concentrated and purified by flash chromatography.

Examples of obtaining the compounds of formula VIII



VIII

Intermediate No.	R ¹	R ²	R
A	Me		H
B	Me		Cl

5 are provided below:

Compound VIII A

4-(4-chloro-*N*-methylanilino)quinoline.

A solution of 4-chloroquinoline (5 mmol) and of 4-chloro-*N*-methylaniline (10 mmol) in glacial acetic acid (15 ml) was
 10 heated with reflux for 3 h under a stream of argon. After cooling, the solution was basified with a 10% NaOH solution until pH = 10 and the resulting suspension was concentrated in a rotary evaporator and purified by means of flash chromatography (9:1, CH₂Cl₂:MeOH) to give the target molecule as a yellowish
 15 syrup (97%). ¹H-NMR (400 MHz, CDCl₃): δ 8.10 (d, *J* = 8.5, 1H, H-2_{quin}); 7.70 (d, *J* = 8.5, 1H, H-5_{quin}); 7.65 (t, *J* = 7.9, 1H, H-7_{quin}); 7.38 (t, *J* = 8.5, 1H, H-6_{quin}); 7.35 (d, *J* = 7.9, 1H, H-8_{quin}); 7.17 (d, *J* = 8.9, 2H, H-3,5_{anil}); 7.14 (d, *J* = 8.5, 1H, H-3_{quin}); 6.76 (d, *J* = 8.9, 2H, H-2,6_{anil}); 3.45 (s, 6H, Me). ¹³C-NMR
 20 (100 MHz, CDCl₃): δ 153.37 (C-4_{quin}); 151.16 (C-2_{quin}); 150.01 (C-1_{anil}); 148.17 (C-8_{aquin}); 135.02 (C-4_{anil}); 130.07 (C-7_{quin}); 129.52 (C-6_{quin}); 129.29 (C-3,5_{anil}); 126.26 (C-4_{aquin}); 126.07 (C-5_{quin}); 124.40 (C-8_{quin}); 119.79 (C-2,6_{anil}); 115.08 (C-3_{quin}); 41.75 (Me). HRMS (*m/z*): Calculated for C₁₆H₁₃N₂Cl [(*M* + *H*)]⁺ 269.0845. Found:
 25 269.0845. Analysis for C₁₆H₁₃N₂Cl. Calculated: C 71.51; H 4.88; N 10.42%. Found: C 71.60; H 4.71; N 10.33%.

Compound VIII B

7-chloro-4-(4-chloro-*N*-methylanilino)quinoline.

A solution of 4,7-dichloroquinoline (5 mmol) and of 4-chloro-*N*-methylaniline (10 mmol) in glacial acetic acid (15 ml) was heated with reflux for 3 h under a stream of argon. After cooling, the solution was basified with a 10% NaOH solution until pH = 10 and the resulting suspension was concentrated in a rotary evaporator and purified by means of flash chromatography (9:1, CH₂Cl₂:MeOH) to give the intermediate II as a yellowish syrup (59%). ¹H-NMR (300 MHz, CH₃OD): δ 8.66 (d, *J* = 7.1, 1H, H-2_{quin}); 7.94 (d, *J* = 2.0, 1H, H-8_{quin}); 7.53 (d, *J* = 8.8, 2H, H-3,5_{anil}); 7.41-7.37 (m, 2H, H-5,6_{quin}); 7.47 (d, *J* = 8.8, 2H, H-2,6_{anil}); 7.32 (d, *J* = 7.1, 2H, H-3_{quin}); 3.76 (s, 3H, Me). ¹³C-NMR (75 MHz, CH₃OD): δ 159.86 (C-4_{quin}); 147.63 (C-7_{quin}); 143.86 (C-2_{quin}); 141.46 (C-1_{anil}); 140.56 (C-8a_{quin}); 135.02 (C-4_{anil}); 132.01 (C-3,5_{anil}); 129.92 (C-6_{quin}); 128.58 (C-2,6_{anil}); 127.98 (C-5_{quin}); 120.56 (C-8_{quin}); 118.71 (C-4a_{quin}); 107.38 (C-3_{quin}); 45.74 (Me). HRMS (*m/z*): Calculated for C₁₆H₁₂N₂Cl₂ [(*M* + *H*)]⁺ 303.0456. Found: 303.0456. Analysis for C₁₆H₁₂N₂Cl₂. Calculated: C 63.38; H 3.99; N 9.24%. Found: C 63.46; H 3.71; N 9.17%.

EX VIVO ASSAYS OF THE ACTIVITY OF HUMAN CHOK

Recombinant choline kinase expressed in *E. coli* in the buffer assay (100 mM Tris-HCl pH 8.0, 100 mM MgCl₂, 10 mM ATP and 200 μM of choline in the presence of methyl[¹⁴C]-choline chloride (50-60 μCi/mmol)) was used for the *ex vivo* assays. The reactions were carried out at 37° C for 30 minutes and were stopped with trichloroacetic acid cooled with ice to a final concentration of 16%. The samples were washed with diethyl ether saturated with water and were lyophilized. The hydrophilic choline derivatives were resolved in thin layer chromatography plates according to a described process [Ramírez, A., Penalva, V., Lucas, L., Lacal, J.C. *Oncogene* 21, 937-946 (2002)].

These assays were carried out with compounds 1-10 of the invention as well as with the compounds EC1-EC6, known compounds of the state of the art, specifically in patent ES 2 117 950. The results are summarized in table II.

CELL PROLIFERATION ASSAYS

The HT-29 cells were seeded in 24-well plates (35 x 10³

cells/well) and were incubated for 24 hours. Then the cells were treated with different concentrations of ChoK inhibitors in the usual culture medium. Three days later, the wells were aspirated and both fresh medium and more amount of drug were added, and
5 the cells were maintained for three more days. The quantification of the remaining cells in each well was carried out by means of the Crystal Violet method [Gillies, R. J., Didier, N., Denton, M. *Anal. Biochem.* **159**, 109-113 (1986)], with some modifications [Hernández-Alcoceba, R., Saniger, L., Campos,
10 J., Núñez, M. C., Khaless, F., Gallo, M. Á., Espinosa, A., Lacal, J. C. *Oncogene*, **15**, 2289-2301 (1997)]. Briefly, the cells were washed with TD buffer and fixed with 1% glutaraldehyde for 15 minutes. After washing again with TD, the cell nuclei were coloured with 0.1% Crystal Violet for at least 30 minutes and
15 washed three times with distilled water. The adsorbed colouring was resuspended in 10% acetic acid and the absorbance at 595 nm was determined in a spectrometer. The results obtained are summarized in the form of an IC_{50} value, i.e. the concentration of the compound required to produce 50% inhibition. This value
20 is determined by iterative curve adjustment. Two values are determined for each point of the curve, the experiment was repeated two or three times and the mean values were estimated. In the few cases in which the two values differed more than 50%, a third experiment was carried out to determine the real value.
25 The IC_{50} value as a potency measurement is used to relate the biological activity of the compounds with their chemical structure.

These assays were carried out with compounds 1-10 of the invention as well as with compounds EC1-EC6, known compounds of
30 the state of the art, specifically in patent ES 2 117 950. The results are summarized in table II.

TOXICITY ASSAYS















Toxicity assays were carried out with 1 month old Balb C mice weighing about 25-30 grams at the start of the experiment.
35 The mice were inoculated with different amounts of each compound in a range of 0.1 mg/kg up to 25 mg/kg, in daily doses for 5 consecutive days. After the five doses, the mice were left to

rest for 9 days and both the survival and general condition were analyzed, paying special attention to the effects on their coat, behaviour, feeding habits and weight. The doses entailing a 50% mortality were recorded as the corresponding toxicity IC_{50} . The
5 results obtained with the new compounds show a clear improvement of the activity when their toxicity is reduced, measured by their corresponding IC_{50} .

These assays were carried out with compounds 1-10 of the invention as well as with compounds EC1-EC6, known compounds of
10 the state of the art, specifically in patent ES 2 117 950. The results are summarized in table II.

The following table II summarizes the results obtained in the assays which have been carried out.

Table II

No.	Code	R ₃ , R ₄ *	NR ₁ R ₂	A	ex vivo IC ₅₀ (μ M)	HT-29 IC ₅₀ (μ M)	toxicity IC ₅₀ (mg/Kg)
1	ACG560B	H, H			5.7	3.3	17.5
2	ACG416B	H, H			0.42	2.2	13.6
3	ACG548B	H, H			1.9	1.9	20
4	ACG604A	H, H			2.6	1.8	16.7
EC1	ACG516B	H, H			5.8	2.6	12.5
EC2	ACG492A	H, H			1.9	1.6	11.5
5	RSM964A	-(CH=CH) ₂ -			1.3	1.6	>25

6	RSM820C	$\text{-C}^5\text{H}=\text{C}^6\text{H}-$ $\text{C}^7\text{Cl}=\text{C}^8\text{H}-$			5.70	1.90	>20
EC3	RSM856B	$\text{-C}^5\text{H}=\text{C}^6\text{H}-$ $\text{C}^7\text{Cl}=\text{C}^8\text{H}-$	-NMe ₂		9.60	0.70	2.9
EC4	RSM1076A	$\text{-C}^5\text{H}=\text{C}^6\text{H}-$ $\text{C}^7\text{Cl}=\text{C}^8\text{H}-$			1.20	0.40	10
7	RSM932A	$\text{-(CH=CH)}_2\text{-}$			2.0	1.2	12.5
8	RSM824B	$\text{-C}^5\text{H}=\text{C}^6\text{H}-$ $\text{C}^7\text{Cl}=\text{C}^8\text{H}-$			11.4	1.2	15
9	RSM936A	$\text{-(CH=CH)}_2\text{-}$			4.8	0.7	16.7
10	RSM828B	$\text{-C}^5\text{H}=\text{C}^6\text{H}-$ $\text{C}^7\text{Cl}=\text{C}^8\text{H}-$			5.70	0.80	12.5
EC5	RSM1084A	$\text{-C}^5\text{H}=\text{C}^6\text{H}-$ $\text{C}^7\text{Cl}=\text{C}^8\text{H}-$			1.00	0.20	7.5
EC6	JC/947A	H, H	-NMe ₂		22	2.5	0.3

* R₃ and R₄ can mean either each one is hydrogen or both form a single radical.

It can be observed from the data of table II that the compounds of the present invention have a considerably lower toxicity than the compounds of patent ES 2 117 950, whereas they
5 maintain similar or greater values of antiproliferative activity against cells derived from culture tumors and of *in vivo* antitumor activity against human tumors inoculated in immunodepressed mice.